Modified Oswestry Low Back Pain Disability Questionnaire^a

☐ I am in bed most of the time and have to crawl to

the toilet.

This questionnaire has been designed to give your therapist information as to how your back pain has affected your ability to manage in everyday life. Please answer every question by placing a mark in the **one** box that best describes your condition today. We realize you may feel that two of the statements may describe your condition, but **please mark only the box that most closely describes your current condition.**

Pai	in Intensity	Sit	ting
	I can tolerate the pain I have without having to use		I can sit in any chair as long as I like.
	pain medication.		I can only sit in my favorite chair as long as I like.
	The pain is bad, but I can manage without having		Pain prevents me from sitting for more than 1 hour.
	to take pain medication.		Pain prevents me from sitting for more than
	Pain medication provides me with complete relief		1/2 hour.
	from pain.		Pain prevents me from sitting for more than
	Pain medication provides me with moderate relief		10 minutes.
	from pain.		Pain prevents me from sitting at all.
	Pain medication provides me with little relief		
	from pain.	Sta	anding
	Pain medication has no effect on my pain.		I can stand as long as I want without increased pain.
	7 1		I can stand as long as I want, but it increases
Peı	rsonal Care (e.g., Washing, Dressing)		my pain.
	I can take care of myself normally without causing		Pain prevents me from standing for more than
	increased pain.		1 hour.
	I can take care of myself normally, but it increases		Pain prevents me from standing for more than
	my pain.		1/2 hour.
	It is painful to take care of myself, and I am slow		Pain prevents me from standing for more than
	and careful.		10 minutes.
	I need help, but I am able to manage most of my		Pain prevents me from standing at all.
	personal care.		, I
	I need help every day in most aspects of my care.	Sle	eping
	I do not get dressed, I wash with difficulty, and I		Pain does not prevent me from sleeping well.
	stay in bed.		I can sleep well only by using pain medication.
			Even when I take medication, I sleep less than
Lif	ting		6 hours.
	I can lift heavy weights without increased pain.		Even when I take medication, I sleep less than
	I can lift heavy weights, but it causes increased pain.		4 hours.
	Pain prevents me from lifting heavy weights off		Even when I take medication, I sleep less than
	the floor, but I can manage if the weights are		2 hours.
	conveniently positioned (e.g., on a table).		Pain prevents me from sleeping at all.
	Pain prevents me from lifting heavy weights, but		T T G
	I can manage light to medium weights if they are	Soc	cial Life
	conveniently positioned.		My social life is normal and does not increase
	I can lift only very light weights.		my pain.
	I cannot lift or carry anything at all.		My social life is normal, but it increases my level
			of pain.
Wa	alking		Pain prevents me from participating in more
	Pain does not prevent me from walking any distance.		energetic activities (e.g., sports, dancing).
	Pain prevents me from walking more than 1 mile.		Pain prevents me from going out very often.
	(1 mile = 1.6 km).		Pain has restricted my social life to my home.
	Pain prevents me from walking more than 1/2 mile.		I have hardly any social life because of my pain.
	Pain prevents me from walking more than 1/4 mile.	_	
	I can walk only with crutches or a cane.		

Please complete questionnaire on other side.

My normal homemaking / job activities do not cause pain.
cause pain.
☐ My normal homemaking / job activities increase
my pain, but I can still perform all that is required
of me.
☐ I can perform most of my homemaking / job
 duties, but pain prevents me from performing more physically stressful activities (e.g., lifting, vacuuming). Pain prevents me from doing anything but light duties. Pain prevents me from doing even light duties. Pain prevents me from performing any job or homemaking chores.
first statement is marked the section score = 0, if the last he score is calculated as follows:
he score is calculated as follows: % ted:
he score is calculated as follows:

Source: Fritz JM, Irrgang JJ. A comparison of a modified Oswestry Low Back Pain Disability Questionnaire and the Quebec Back Pain Disability Scale. *Physical Therapy*. 2001;81:776-788.

^aModified by Fritz & Irrgang with permission of The Chartered Society of Physiotherapy, from Fairbanks JCT, Couper J, Davies JB, et al. The Oswestry Low Back Pain Disability Questionnaire. *Physiotherapy*. 1980;66:271-273.